



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GEOMETRY.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

52. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pennsylvania.

If the center of a rolling ellipse move in a horizontal line, determine the surface on which the ellipse rolls.

Solution by G. B. M. ZERE, A. M., Ph. D., Professor of Mathematics and Applied Science, Texarkana College, Texarkana, Arkansas-Texas.

Let BPA be a quadrant of the ellipse semi-axes AC , and BC , O the position of the center when BC coincides with OY , and $\angle BCP = \theta$. Then

$$PC = y = \frac{ab}{\sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta}} = \frac{b}{\sqrt{1 - e^2 \sin^2 \theta}}.$$

\therefore The ellipse rolls on the inner surface of the cylinder

$$y^2 + z^2 = \frac{b^2}{1 - e^2 \sin^2 \theta}.$$

When $e=0$, this becomes $y^2 + z^2 = b^2$.

To find the abscissa of the point of contact, we have, since $\text{arc } PB = \text{arc } PG$,

$$ds = \sqrt{r^2 d\theta^2 + dr^2} = \sqrt{y^2 d\theta^2 + dy^2} \text{ since } PC = r = y;$$

$$\text{also } ds = \sqrt{dx^2 + dy^2}.$$

$$\therefore \sqrt{dx^2 + dy^2} = \sqrt{y^2 d\theta^2 + dy^2}.$$

$$\therefore dx = y d\theta, \text{ or } x = \int y d\theta = \int \frac{bd\theta}{\sqrt{1 - e^2 \sin^2 \theta}} = bF(e, \theta).$$

When $e=0$, $x = b\theta$.

[No other solution of this problem was received. EDITOR.]

53. Proposed by B. F. FINKEL, A. M., Professor of Mathematics and Physics in Drury College, Springfield, Missouri.

A pole, a certain length of whose top is painted white, is standing on the side of a hill. A person at A observes that the white part of the pole subtends an angle equal to α

